

黄皮根和茎中一个新的木脂体*

宋卫武^{1,2}, 曾广智¹, 彭文文¹, 谭宁华^{1**}

(1 中国科学院昆明植物研究所 植物化学与西部植物资源持续利用国家重点实验室, 云南 昆明 650201;

2 中国科学院大学, 北京 100049)

摘要: 从黄皮 (*Clausena lansium*) 根和茎中分离得到 8 个化合物, 其中化合物 claulignan (**1**) 为一个新的木脂体类化合物, 通过 1D-、2D-NMR 和高分辨质谱确定其结构。化合物 **2-8** 首次从黄皮属 (*Clausena*) 植物中分离得到。活性筛选结果表明, 化合物 **1** 和 **6** 显示一定的细胞毒活性, 对人宫颈癌 HeLa 细胞株的 IC_{50} 分别为 25.03 和 53.99 μM ; 化合物 **3** 具有一定的抗氧化活性, 在 DPPH 实验中的 EC_{50} 为 268.96 $\text{g} \cdot \text{kg}^{-1}$ 。

关键词: 黄皮; 木脂体; 细胞毒; DPPH; Claulignan

中图分类号: Q 946

文献标识码: A

文章编号: 2095-0845(2014)04-545-06

A New Cytotoxic Oxynolignan from the Roots and Stems of *Clausena lansium* (Rutaceae)*SONG Wei-Wu^{1,2}, ZENG Guang-Zhi¹, PENG Wen-Wen¹, TAN Ning-Hua^{1**}

(1 State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China; 2 University of Chinese Academy of Sciences, Beijing 100049, China)

Abstract: A new oxynolignan, claulignan (**1**), together with seven known compounds (**2-8**), were isolated from the methanol extract of the roots and stems of *Clausena lansium*. Their structures were elucidated on the basis of 1D- and 2D-NMR experiments and mass spectrometry. Compounds **2-8** were isolated from the genus *Clausena* for the first time. Compounds **1** and **6** showed cytotoxicity against human cervical cancer HeLa cell line with the IC_{50} values of 25.03 and 53.99 μM , respectively. Compound **3** exhibited antioxidant activity in the DPPH assay with the EC_{50} value of 268.96 $\text{g} \cdot \text{kg}^{-1}$.

Key words: *Clausena lansium*; Oxynolignan; Cytotoxicity; DPPH; Claulignan

Clausena belongs to the family Rutaceae and includes about 10 species in China. *Clausena lansium* (Lour.) Skeels is a fruit tree and widely distributes in southern China (Huang, 1997). Its fruits have been used for treating indigestion, cold, cough, and stomach pain; its seeds for treating acute, chronic gastrointestinal inflammation, and ulcer; and its leaves and roots for the treatment of

cough, asthma, dermatological disease, viral hepatitis, and gastro-intestinal disease (Shen *et al.*, 2012). Previous phytochemical investigation revealed that *C. lansium* contained flavonoid glycosides, coumarins, cabazole alkaloids, amides, and terpenoids with cytotoxic and neuroprotective activities (Ito *et al.*, 1998; Lakshmi *et al.*, 1989; Liu *et al.*, 2012; Maneerat *et al.*, 2012; Yang *et al.*, 1988; Zhao *et*

* Funding: Found of the National New Drug Innovation Major Project of China (2011ZX09307-002-02), the National Natural Science Foundation of China (U1032602 and 91013002), the National Basic Research Program of China (2009CB522300), the Fund of Chinese Academy of Sciences (Hundred Talents Program), and the Natural Science Foundation of Yunnan Province (2012GA003)

** Author for correspondence; E-mail: nhtan@mail.kib.ac.cn

Received date: 2013-09-16, Accepted date: 2013-10-10

作者简介: 宋卫武 (1983-) 男, 在读博士研究生, 主要从事天然药物化学研究。

al., 2010). In order to discover bioactive constituents, the methanol extract of the roots and stems of *C. lansium* was studied. Herein we report the isolation and structural elucidation of one new oxyneolignan and 7 known compounds from this extract with cytotoxic and antioxidant activities.

1 Results and discussion

Claulignan (**1**) was obtained as a brown powder and its molecular formula was deduced as $C_{20}H_{22}O_7$ on the basis of HR-EI-MS at m/z 374.1378 $[M]^+$ (calcd for $C_{20}H_{22}O_7$, 374.1366). The ^{13}C -NMR spectrum of **1** (Table 1) displayed ten carbon signals, including six aromatic carbons (δ_C 152.3, 148.3, 130.6, 124.0, 115.4 and 111.5), two methylene carbons (δ_C 58.7 and 41.5), one keto-carbon (δ_C 198.2) and one methoxy carbon (δ_C 56.2). In the 1H -NMR spectrum of **1** (Table 1), three aromatic proton signals at δ_H 7.55 (d, $J=1.5$ Hz), 6.92 (d, $J=8.2$ Hz) and 7.59 (dd, $J=8.2, 1.5$ Hz) were observed. These data indicated that **1** might be an oxyneolignan with the same unit of 3, 4-disubstituted phenylpropanoid. Position of the keto-group was determined by the HMBC correlations from H-2, H-6, H-8 and H-9 to C-7, and the COSY correlation between H-8 and H-9 (Fig. 1). The HMBC correlation from OCH_3 to C-3 and the ROESY correlation between H-2 and OCH_3 suggested that the methoxy group linked at C-3 and the two same phenylpropanoid units linked at C-4 (Fig. 2). Thus, the structure of **1** was established.

Seven known compounds were identified to be clemaphenol A (**2**) (Song *et al.*, 2010), syringaresinol (**3**) (Ouyang *et al.*, 2007), ficusol (**4**) (Li and Kuo, 1998), berfussinol (**5**) (Kim *et al.*, 2011), 4-hydroxy-2-methoxycinnamaldehyde (**6**) (Xu and Wang, 2011), 3-methoxy-4-hydroxybenzaldehyde (**7**) (Yang *et al.*, 2007) and 4-hydroxybenzaldehyde (**8**) (Feng *et al.*, 2008) (Fig. 3). Compounds **2–8** were isolated from the genus *Clausea* for the first time. Compounds **1** and **6** showed cytotoxicity against human cervical cancer HeLa cell

line with the IC_{50} values of 25.03 and 53.99 μM , respectively. Compound **3** exhibited antioxidant activity in the DPPH assay with the EC_{50} value of 268.96 $g \cdot kg^{-1}$.

Table 1 1H and ^{13}C NMR data for compound **1**
in acetone- d_6 (δ in ppm, J in Hz)

NO.	δ_C^a	δ_H^b
1, 1'	130.6 (s)	
2, 2'	111.5 (d)	7.55 (d, 1.5)
3, 3'	148.3 (s)	
4, 4'	152.3 (s)	
5, 5'	115.4 (d)	6.92 (d, 8.2)
6, 6'	124.0 (d)	7.59 (dd, 8.2, 1.5)
7, 7'	198.2 (s)	
8, 8'	41.5 (t)	3.16 (m)
9, 9'	58.7 (t)	3.91 (overlapped)
OH		8.70 (s)
OMe	56.2 (q)	3.90 (s)

a) 600 MHz; b) 100 MHz




Fig. 1 Structure of compound **1**




Fig. 2 The key HMBC (→) and 1H , 1H -COSY (—) correlations for compound **1**

2 Experimental

General experimental procedures Optical rotation was measured on a Jasco P-1020 polarimeter. UV spectrum was obtained from Shimadzu UV-2401PC spectrophotometer. IR spectrum was recorded on a BRUKER Tensor 27 FT-IR spectrometer with KBr pellets. The 1D- and 2D-NMR spectra were




Fig. 3 Structures of compounds 2–8

obtained from Bruker AM-400 (1H : 400 MHz, ^{13}C : 100 MHz) or Bruker DRX-500 (1H : 500 MHz, ^{13}C : 150 MHz) or Bruker AV-III 600 (1H : 600 MHz, ^{13}C : 150 MHz) spectrometer in acetone- d_6 , CD_3OD or pyridine- d_5 at the room temperature. ESI-MS was determined on a Waters Xevo TQ-S or Bruker HCT/Esquire. EI-MS and HR-EI-MS were determined on a Waters AutoSpec Premier P776. Silica gel (200–300 mesh; Qingdao Yuminyuan Silica Reagent Factory, Qingdao, P. R. China), RP-18 (40–60 μm , Merck, Darmstadt, Germany), and MCI gel (CHP-20P, 70–150 μm , Mitsubishi Chemical Corporation, Japan) were used for column chromatography (CC). Fractions were monitored by TLC (Silica gel GF₂₅₄ Plates, Qingdao Yuminyuan Silica Reagent Factory, Qingdao, P. R. China) and spots were visualized by UV light (254 and 365 nm), and by spraying with 5% aq. H_2SO_4 soln. followed by heating. MPLC, equipped with a UV detector (EZ Purifier 100/200, Lisure Science (Suzhou) Co., LTD. P. R. China) and a RP-18 column (3.5×25 or 4.5×40 cm) or a MCI column (4.5×40 cm) at a flow rate of 25 mL·min⁻¹, and preparative reversed-phase HPLC, Agilent 1100 apparatus equipped with a UV

detector and a SunFire OBD (Waters, 1.9×25 cm, 5 μm) column at a flow rate of 10 mL·min⁻¹ were used for purification.

Plant material The roots and stems of *Clausena lansium* were collected from Hekou, Honghe Hani & Yi Autonomous Prefecture, Yunnan province, P. R. China, in September 2010 and identified by Prof. Yu-Min Shui, Kunming Institute of Botany, Chinese Academy of Sciences. A voucher specimen (No. 0599043) was deposited at the Herbarium of Kunming Institute of Botany, Chinese Academy of Sciences.

Extraction and isolation Air dried, powdered roots and stems of *C. lansium* (27 kg) were extracted and refluxed with MeOH for 3 times each 4 hours (MeOH, 50 L×3). The extract was evaporated under reduced pressure to yield a dark brown residue (900 g). The residue was suspended in MeOH/ H_2O (7:3, 3 000 mL) and then partitioned with EtOAc (3×2 000 mL). After removing solvent, the EtOAc-soluble part (406 g) was fractionated by silica gel (200–300 mesh) column chromatograph (CC) and eluted with $CHCl_3$ -MeOH (30:1–4:1) to afford 6 fractions, Fractions 1–6 (Fr. 1–6), on the basis of

TLC analysis. Fr. 3 (25 g) was chromatographed over MPLC with MCI (MeOH/H₂O 10:90→100:0) and silica gel (petroleum ether/acetone 15:1→7:3) successively to afford **6** (15 mg) and **7** (10 mg). Fr. 4 (17.9 g) was firstly subjected to MPLC with MCI column (MeOH/H₂O 10:90→60:40) to remove pigment and then to silica gel CC (petroleum ether/acetone 9:1) and preparative HPLC (acetonitrile/H₂O 55:45) to afford **2** (26 mg). MPLC chromatograph of Fr. 5 (6.9 g) over MCI (MeOH/H₂O 5:95→60:40) afforded 7 sub-fractions: Fr. 5-1-Fr. 5-7. Fr. 5-1 was repeatedly chromatographed over silica gel (petroleum ether/acetone 9:1) and then subjected to preparative HPLC (acetonitrile/H₂O 55:45) to afford **1** (22 mg), **3** (280 mg), **4** (94 mg) and **5** (35 mg). Fr. 6 (77 g) was subjected to silica gel CC (CHCl₃/acetone 15:1→7:3), silica gel CC (petroleum ether/acetone 5:1), MPLC with MCI (MeOH/H₂O 10:90→60:40), MPLC with RP-18 (MeOH/H₂O 5:95→70:30) and silica gel CC (petroleum ether/EtOAc 5:1), subsequently to afford **8** (21 mg).

Claulignan (1) was obtained as brown powder, C₂₀H₂₂O₇; $[\alpha]_D^{21.0} = -12.93$ ($c = 0.10$, MeOH); UV (MeOH) λ_{\max} (log ϵ): 205 (4.48), 228 (4.48), 276 (4.30), 303 (4.23) nm; IR (KBr) ν_{\max} (cm⁻¹): 3423, 2960, 1665, 1591, 1517, 1464, 1427, 1385, 1279, 1197, 1167, 1136, 1032; ¹H NMR (acetone-d₆, 600 MHz) and ¹³C NMR (acetone-d₆, 100 MHz): see Table 1; ESI-MS (positive) m/z : 397 [M+Na]⁺, 771 [2M+Na]⁺; HR-EI-MS m/z : 374.1378 [M]⁺ (calcd. for C₂₀H₂₂O₇⁺, 374.1366).

Clemaphenol A (2) was obtained as colorless oil, C₂₀H₂₂O₆; $[\alpha]_D^{23.7} = +0.16$ ($c = 0.42$, MeOH); ¹H NMR (400 MHz, acetone-d₆): δ_H 6.98 (2H, d, $J = 1.3$ Hz, H-2, H-2'), 6.83 (1H, dd, $J = 8.1, 1.3$ Hz, H-6, H-6'), 6.78 (2H, d, $J = 8.1$ Hz, H-5, H-5'), 4.65 (2H, d, $J = 3.8$ Hz, H-7, H-7'), 4.20 (2H, m, H_a-9, H_a-9'), 3.83 (6H, s, OCH₃), 3.80 (2H, dd, $J = 9.2, 3.8$ Hz, H_b-9, H_b-9'), 3.10 (2H, m, H-8,

H-8'); ¹³C NMR (100 MHz, acetone-d₆): δ_C 148.2 (s, C-4, C-4'), 146.8 (s, C-3, C-3'), 134.1 (s, C-1, C-1'), 119.6 (d, C-6, C-6'), 115.5 (d, C-5, C-5'), 110.5 (d, C-2, C-2'), 86.6 (d, C-7, C-7'), 72.1 (t, C-9, C-9'), 56.2 (q, OCH₃), 55.2 (d, C-8, C-8'); ESI-MS (positive) m/z : 381 [M+Na]⁺, 739 [2M+Na]⁺. (Song *et al.*, 2010)

Syringaresinol (3) was obtained as colorless oil, C₂₂H₂₆O₈; $[\alpha]_D^{21.5} = -6.7$ ($c = 0.20$, MeOH); ¹H NMR (400 MHz, CD₃OD): δ_H 6.51 (4H, s, H-2, H-2', H-6, H-6'), 4.54 (2H, d, $J = 3.8$ Hz, H-7, H-7'), 4.09 (2H, m, H_a-9, H_a-9'), 3.71 (2H, dd, $J = 9.3, 2.8$ Hz, H_b-9, H_b-9'), 3.67 (12H, s, OCH₃), 2.97 (2H, br. s, H-8, H-8'); ¹³C NMR (100 MHz, CD₃OD): δ_C 149.1 (s, C-3, C-3', C-5, C-5'), 135.9 (s, C-4, C-4'), 133.0 (s, C-1, C-1'), 104.3 (d, C-2, C-2', C-6, C-6'), 87.4 (d, C-7, C-7'), 72.6 (t, C-9, C-9'), 56.7 (q, OCH₃), 55.3 (d, C-8, C-8'); ESI-MS (negative) m/z : 417 [M-H]⁻, 418 [M]⁻. (Ouyang *et al.*, 2007)

Ficusol (4) was obtained as colorless oil, C₁₁H₁₄O₅; $[\alpha]_D^{21.7} = -5.5$ ($c = 0.38$, MeOH); ¹H NMR (400 MHz, acetone-d₆): δ_H 6.92 (1H, d, $J = 0.8$ Hz, H-2'), 6.77 (2H, overlapped, H-5', H-6'), 4.07 (1H, m, H_a-3), 3.82 (3H, s, 3'-OCH₃), 3.74 (1H, overlapped, H_b-3), 3.68 (1H, overlapped, H-2), 3.63 (3H, s, 1-OCH₃); ¹³C NMR (100 MHz, acetone-d₆): δ_C 174.1 (s, C-1), 148.3 (s, C-3'), 146.8 (s, C-4'), 128.6 (s, C-1'), 121.5 (d, C-6'), 115.8 (d, C-5'), 112.5 (d, C-2'), 65.2 (t, C-3), 56.2 (q, 3'-OCH₃), 54.9 (d, C-2), 52.0 (q, 1-OCH₃); ESI-MS (positive) m/z : 249 [M+Na]⁺. (Li and Kuo, 1998)

Berfussinol (5) was obtained as white solid, C₁₂H₁₆O₆; $[\alpha]_D^{21.7} = -5.8$ ($c = 0.15$, MeOH); ¹H NMR (400 MHz, acetone-d₆): δ_H 6.60 (2H, s, H-2', H-6'), 4.07 (1H, m, H_a-3), 3.79 (6H, s, 3'-OCH₃, 5'-OCH₃), 3.71 (1H, overlapped,

H_b-3), 3.70 (1H, overlapped, H-2), 3.63 (3H, s, 1-OCH₃); ¹³C NMR (100 MHz, acetone-d₆): δ_C 174.0 (s, C-1), 148.6 (s, C-3', C-5'), 136.2 (s, C-4'), 127.5 (s, C-1'), 106.5 (d, C-2', C-6'), 65.2 (t, C-3), 56.6 (q, 3'-OCH₃, 5'-OCH₃), 55.1 (d, C-2), 51.9 (q, 1-OCH₃). ESI-MS (positive) *m/z*: 279 [M+Na]⁺, 535 [2M+Na]⁺. (Kim *et al.*, 2011)

4-Hydroxy-2-methoxycinnamaldehyde (6)

was obtained as yellow oil, C₁₀H₁₀O₃; ¹H NMR (600 MHz, acetone-d₆): δ_H 9.63 (1H, d, *J*=7.8 Hz, H-9), 8.43 (1H, s, OH), 7.57 (1H, d, *J*=15.8 Hz, H-7), 7.39 (1H, d, *J*=2.0 Hz, H-3), 7.21 (1H, dd, *J*=8.0, 2.0 Hz, H-5), 6.91 (1H, d, *J*=8.0 Hz, H-6), 6.67 (1H, dd, *J*=15.8, 7.8 Hz, H-8), 3.92 (3H, s, OCH₃); ¹³C NMR (150 MHz, acetone-d₆): δ_C 194.0 (d, C-9), 154.2 (d, C-7), 150.9 (s, C-4), 148.9 (s, C-2), 127.4 (s, C-1), 127.0 (d, C-8), 124.9 (d, C-6), 116.2 (d, C-5), 111.4 (d, C-3), 56.3 (q, OCH₃); ESI-MS (negative) *m/z*: 177 [M-H]⁻, 178 [M]⁻, 355 [2M-H]⁻. (Xu and Wang, 2011)

3-Methoxy-4-hydroxybenzaldehyde (7) was obtained as white powder, C₈H₈O₃; ¹H NMR (600 MHz, acetone-d₆): δ_H 9.83 (1H, s, CHO), 8.77 (1H, s, OH), 7.46 (1H, dd, *J*=7.9, 1.8 Hz, H-6), 7.44 (1H, d, *J*=1.8 Hz, H-2), 7.01 (1H, d, *J*=7.9 Hz, H-5), 3.93 (3H, s, OCH₃); ¹³C NMR (150 MHz, acetone-d₆): δ_C 191.2 (d, CHO), 153.6 (s, C-4), 149.0 (s, C-3), 130.8 (s, C-1), 127.10 (d, C-6), 116.0 (d, C-5), 110.8 (d, C-2), 56.3 (q, OCH₃); ESI-MS (negative) *m/z*: 151 [M-H]⁻, 152 [M]⁻. (Yang *et al.*, 2007)

4-Hydroxybenzaldehyde (8) was obtained as colorless crystal, C₇H₆O₂; ¹H NMR (400 MHz, pyridine-d₅): δ_H 10.00 (1H, s, CHO), 7.94 (2H, d, *J*=8.0 Hz, H-2, H-6), 7.21 (2H, d, *J*=8.0 Hz, H-3, H-5); ¹³C NMR (100 MHz, pyridine-d₅): δ_C 190.8 (d, CHO), 165.0 (s, C-4), 132.8 (d, C-2, C-6), 129.9 (s, C-1), 116.8 (d, C-3, C-5); ESI-MS (positive) *m/z*: 123 [M+

H]⁺, 145 [M+Na]⁺. (Feng *et al.*, 2008)

Cytotoxic Assay All compounds were tested for their cytotoxicity against A549, HeLa and BGC-823 cancer cell lines with the sulforhodamine B assay (He *et al.*, 2011). Cells were cultured in RPMI 1640 medium (Gibco) supplemented with 10% FBS (Tianjin Haoyang) at 37 °C and 5% CO₂. After seeded in 96-well microtiter plates for 24 h, cells were treated with serial dilutions of the tested compounds with the maximum concentration of 20 μg · mL⁻¹. Each compound was initially dissolved in DMSO and further diluted by the medium to produce different concentrations. After incubation for 48 or 72 h, cells were fixed with ice-cold 50% trichloroacetic acid at 4 °C for 1 h. With staining for 15 min by 0.4% SRB (Sigma) in 1% glacial acetic acid, excessive dye was removed by washing with 1% glacial acetic acid. Finally, SRB was re-suspended in 10 mmol · L⁻¹ Tris buffer and the absorbance was measured at 560 nm with a plate reader (Molecular Devices, SPECTRA MAX 340). The result was expressed as IC₅₀ value, which is the concentration that causes 50% growth inhibition. Taxol was used as a reference compound.

DPPH assay The antioxidant activity of all compounds was determined using 1, 1-diphenyl-2-picrylhydrazyl (DPPH) (TCI) method according to the procedure described by us (Adebayo *et al.*, 2010). Each compound solution in dimethyl sulphoxide (DMSO, 5 μL) was added to 195 μL of 0.025 g · L⁻¹ DPPH methanol solution. Absorbance at 515 nm was taken after 30 minutes by incubated at room temperature in the microplate reader (Molecular Devices, SPECTRA MAX 340). If the absorbance decreased over 50% by compared with the DMSO control at the concentration of 20 μg · mL⁻¹, at least four serially diluted concentrations of tested compounds were prepared to evaluate their antioxidant activity using the above procedure. The result was expressed as EC₅₀ value, which is the amount to decrease the initial DPPH concentration by 50%. Vitamin C was used as a reference compound.

Acknowledgement: The authors are grateful to the members of the analytical group from the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, for measuring the $[\alpha]$, IR, UV, NMR and MS spectra.

References:

- Adebayo AH, Tan NH, Akindahunsi AA *et al.*, 2010. Anticancer and antiradical scavenging activity of *Ageratum conyzoides* L. (Asteraceae) [J]. *Pharmacognosy Magazine*, **6**: 62—66
- Feng BM (冯宝民), Yu ZJ (余正江), Duan LX (段礼新) *et al.*, 2008. Chemical constituents of roasted *Sinapis* seeds [J]. *Chinese Traditional and Herbal Drugs* (中草药), **39**: 331—334
- He WJ, Chu HB, Zhang YM *et al.*, 2011. Antimicrobial, cytotoxic lignans and terpenoids from the twigs of *Pseudolarix kaempferi* [J]. *Planta Medica*, **77**: 1924—1931
- Huang CJ (黄成就), 1997. Rutaceae [A]. In: Wu ZY (ed.), *Flora of China* (中国植物志) [M]. Beijing: Science Press, Vol. **43**: 132
- Ito C, Katsuno S, Furukawa H, 1998. Structures of lansiumarin-A, -B, -C, three new furocoumarins from *Clausena lansium* [J]. *Chemical & Pharmaceutical Bulletin*, **46**: 341—343
- Kim KH, Moon E, Choi SU *et al.*, 2011. Biological evaluation of phenolic constituents from the trunk of *Berberis koreana* [J]. *Bioorganic & Medicinal Chemistry Letters*, **21**: 2270—2273
- Lakshmi V, Raj K, Kapil RS, 1989. A triterpene alcohol, lansiol, from *Clausena lansium* [J]. *Phytochemistry*, **28**: 943—945
- Li YC, Kuo YH, 1998. A monoterpenoid and two simple phenols from heartwood of *Ficus microcarpa* [J]. *Phytochemistry*, **49**: 2417—2419
- Liu H, Li CJ, Yang JZ *et al.*, 2012. Carbazole alkaloids from the stems of *Clausena lansium* [J]. *Journal of Natural Products*, **75**: 677—682
- Maneerat W, Ritthiwigrom T, Cheenpracha S *et al.*, 2012. Carbazole alkaloids and coumarins from *Clausena lansium* roots [J]. *Phytochemistry Letters*, **5**: 26—28
- Ouyang MA, Wein YS, Zhang ZK *et al.*, 2007. Inhibitory activity against tobacco mosaic virus (TMV) replication of pinoresinol and syringaresinol lignans and their glycosides from the root of *Rhus javanica* var. *roxburghiana* [J]. *Journal of Agricultural and Food Chemistry*, **55**: 6460—6465
- Shen DY, Chao CH, Chan HH *et al.*, 2012. Bioactive constituents of *Clausena lansium* and a method for discrimination of aldose enantiomers [J]. *Phytochemistry*, **82**: 110—117
- Song WW, Li B, Liu JK, 2010. A new lignan from *Elaeagnus lanceolata* (Elaeagnaceae) [J]. *Acta Botanica Yunnanica* (云南植物研究), **32**: 455—462
- Xu C (徐冲), Wang ZT (王峥涛), 2011. Chemical constituents from roots of *Andrographis paniculata* [J]. *Acta Pharmaceutica Sinica* (药学报), **46**: 317—321
- Yang MH, Chen YY, Huang L, 1988. Three novel cyclic amides from *Clausena lansium* [J]. *Phytochemistry*, **27**: 445—450
- Yang XW (杨秀伟), Han MH (韩美华), Jin YP (靳彦平), 2007. Studies on the chemical constituents from herba anoec-tochili [J]. *Journal of Chinese Medicinal Materials* (中药材), **30**: 797—800
- Zhao Q (赵青), Li C (李创), Yang JZ (杨敬芝) *et al.*, 2010. Chemical constituents of *Clausena lansium* [J]. *China Journal of Chinese Materia Medica* (中国中药杂志), **35**: 997—1000